

PREY SELECTION OF SPIDERS IN THE FIELD

Martin Nyffeler: Zoological Institute, Division of Ecology, University of Berne,
Baltzerstr. 3, CH-3012 Berne, Switzerland

ABSTRACT. In this article, an overview of the general feeding patterns of common agroecosystem spiders is presented. Five groups of web-weavers (Tetragnathidae, Araneidae, Theridiidae, Linyphiidae, Dictynidae) and five groups of hunters (small-sized Oxyopidae, large-sized Oxyopidae, Thomisidae, Salticidae, Lycosidae) are analyzed comparatively (based on 40 prey analyses previously published by various European and US authors). Fewer than 10 insect orders, as well as the order Araneae, make up the bulk of the prey of these spiders. Web-weavers and hunters both basically feed on the same prey orders, but in different proportions. The observed differences reflect in part the very diverse range of life styles and foraging modes exhibited by the various spider groups and, to some extent, differences in prey availability. Web-weavers are almost strictly insectivorous (insects constituting > 99% of total prey). Hunters, however, exhibit a mixed strategy of insectivorous and araneophagic foraging patterns (insects constituting ≈75–90% of total prey). Diet breadth computed with the Inverted Simpson Index was, on average, significantly higher in the hunting spiders than the web spiders. There seems to be a consistent trend of greater diet breadth of the hunters compared to the web-weavers in agroecosystems. Overall, spider individuals of small size (including large percentages of immatures) numerically dominate the faunas of field crops, and these feed primarily on tiny prey (< 4 mm in length).

Information on how prey selection in the field operates is a prerequisite to a quantitative assessment of the spiders' potential as biological control agents in agroecosystems. Prey selection has been defined by Hassell (1978) as follows: "Preference for a particular prey is normally measured in terms of the deviation of the proportion of that prey attacked from the proportion available in the environment." Most authors who studied the prey of spiders failed to record the availability of potential prey in the environment probably due to technical difficulties. Thus, corresponding data on the actual and potential prey are scarce; and, consequently, only a limited number of prey selection studies on spiders following Hassell's approach exist (e.g., Uetz et al. 1978).

Another approach to searching for patterns of prey selection is to analyze a large set of data on the actual prey of different spider groups (with very differing life styles and foraging modes) and to compare the degree to which utilization of the various prey taxa differs. Numerous published field studies on the actual prey of spiders are available for such an investigation (see reviews by Nyffeler 1982; Nentwig 1987; Riechert & Harp 1987; Wise 1993; Nyffeler et al. 1994a, b). In the current investigation, five groups of web-

weavers (Tetragnathidae, Araneidae, Theridiidae, Linyphiidae, Dictynidae) and five groups of hunters (small-sized Oxyopidae [i.e., *Oxyopes salticus*], large-sized Oxyopidae [i.e., *Peucetia viridans*], Thomisidae, Salticidae, Lycosidae), representing nine families, are analyzed comparatively. These selected groups are among the most common spider predators in agroecosystems (Nyffeler et al. 1994b) and, thus, are of particular interest from the point of view of biological control. Descriptions of the life styles and foraging modes of these 10 spider groups are given by Rypstra (1982), Nentwig (1987), Wise (1993), and Nyffeler et al. (1994a, b).

METHODS

For each of the 10 spider groups the relative taxonomic composition of the diets (mean \pm SE of 4 different prey analyses) was assessed (Tables 2, 3). Overall, 40 different prey analyses (based on observational data from 31 published studies [see Table 1]) have been processed. To determine relative feeding specialization, the diet breadth B (= diversity of arthropod orders in the diet) was computed for each spider group by means of the Inverted Simpson Index (see Levins 1968; Colwell & Futuyma 1971) (Table 4). Diet breadth is inversely related to ecological specialization

Table 1.—Field studies used for the assessment of the relative taxonomic composition of the diets of ten spider groups. Habitats: SO = soybean, CO = cotton, PE = peanuts, AA = alfalfa, WW = winter wheat, OA = oats, MA = maize, MM = mown meadow, VE = vegetables, NC = noncrop.

Spider group	Habitat	Area	Author(s)
Tetragnathidae			
<i>Tetragnatha laboriosa</i>	SO	USA	LeSar & Unzicker (1978)
<i>Tetragnatha laboriosa</i>	SO	USA	Culin & Yeargan (1982)
<i>Tetragnatha laboriosa</i>	CO	USA	Nyffeler et al. (1989)
<i>Tetragnatha extensa</i>	WW	Europe	Nyffeler & Benz (1979)
Araneidae			
<i>Acanthepeira stellata</i>	CO	USA	Nyffeler et al. (1989)
<i>Argiope aurantia</i>	CO	USA	Nyffeler et al. (1987a)
<i>Neoscona arabesca</i>	CO	USA	Nyffeler et al. (1989)
<i>Neoscona arabesca</i>	SO	USA	Culin & Yeargan (1982)
Theridiidae			
<i>Latrodectus mactans</i>	CO	USA	Nyffeler et al. (1988a)
<i>Achaearanea riparia</i>	WW	Europe	Nyffeler & Benz (1988a)
<i>Theridion impressum</i>	WW	Europe	Nyffeler (1982)
<i>Theridion impressum</i>	OA	Europe	Nyffeler & Benz (1979)
Linyphiidae			
various Erigoninae	MA	Europe	Alderweireldt (1994)
various Erigoninae	WW	Europe	Sunderland et al. (1986)
various Erigoninae	WW	Europe	Nyffeler & Benz (1988b)
various Erigoninae	MM	Europe	Nyffeler (1982)
Dictynidae			
<i>Dictyna segregata</i>	CO	USA	Nyffeler et al. (1988b)
<i>Dictyna arundinacea</i>	WW	Europe	Heidger & Nentwig (1989)
<i>Dictyna arundinacea</i>	NC	Europe	Heidger & Nentwig (1986)
<i>Dictyna montana</i>	NC	Africa	Nentwig (1987)
Oxyopidae (small-sized)			
<i>Oxyopes salticus</i>	CO	USA	Nyffeler et al. (1987b)
<i>Oxyopes salticus</i>	CO	USA	Nyffeler et al. (1992a)
<i>Oxyopes salticus</i>	CO	USA	Lockley & Young (1987)
<i>Oxyopes salticus</i>	PE	USA	Agnew & Smith (1989)
Oxyopidae (large-sized)			
<i>Peucea viridans</i>	CO	USA	Nyffeler et al. (1987c)
<i>Peucea viridans</i>	CO	USA	Nyffeler et al. (1992a)
<i>Peucea viridans</i>	NC	USA	Turner (1979)
<i>Peucea viridans</i>	NC	USA	Randall (1982)
Thomisidae			
<i>Misumenops</i> spp.	PE	USA	Agnew & Smith (1989)
<i>Misumenops</i> spp.	CO, NC	USA	Dean et al. (1987)
<i>Xysticus emertoni</i>	NC	USA	Morse (1983)
<i>Xysticus</i> spp.	MM	Europe	Nyffeler & Breene (1990a)
Salticidae			
<i>Phidippus audax</i>	CO, NC	USA	Dean et al. (1987)
<i>Phidippus audax</i>	CO, NC	USA	Young (1989)
<i>Phidippus audax</i>	VE	USA	Riechert & Bishop (1990)
<i>Phidippus johnsoni</i>	NC	USA	Jackson (1977)
Lycosidae			
<i>Pardosa ramulosa</i>	AA	USA	Yeargan (1975)
<i>Pardosa</i> spp.	PE	USA	Agnew & Smith (1989)
<i>Pardosa</i> spp.	WW	Europe	Nyffeler & Benz (1988c)
<i>Pardosa amentata</i>	NC	Europe	Hallander (1970)

in part, the diverse range of life styles and foraging modes exhibited by the various spider groups, and to some extent differences in prey availability (see Riechert & Luczak 1982; Nentwig 1987; Nyffeler et al. 1994b).

Web-weavers are almost strictly insectivorous (insects constituting > 99% of total prey) (Table 2). Aggressive encounters among web-weavers occur quite frequently, but rarely result in predation. In a web, the potential victim gets advanced vibrational warning and can flee or be ready to repulse the attacker. During such encounters between web-weavers the inferior individual is usually chased away by its opponent (see Wise 1993). Under conditions of suitable food supply in the form of insects the web-weavers seem to minimize feeding on "dangerous prey" such as spiders. Hunters, however, exhibit a mixed strategy of insectivorous and araneophagic foraging patterns (insects constituting \approx 75–90% of total prey) (Table 3). Field populations of several species of hunters had been found to be in a state of undernourishment (see Nyffeler & Breene 1990b). Thus, araneophagy including cannibalism (as an additional feeding strategy to insectivory) may be crucial in sustaining the hunter populations during periods of food shortage (see Wise 1993). "Eating other spiders appears to be an opportunistic occurrence, a larger or faster individual overpowering another in a chance encounter" (Jackson 1992).

Based on the data presented in Tables 2 and 3, the diet breadth (B) for spiders was computed with the Inverted Simpson Index (Table 4). The highest value was approximately five times higher than the minimum ($B = 1.13$ vs. 5.58), which indicates considerable between-species differences in diet breadth. Evidently the hunters exhibit on average a less specialized feeding behavior (overall mean diet breadth = 4.20 ± 0.20) compared to the web-weavers (overall mean = 2.61 ± 0.22) (Table 4), the difference between the two overall means being statistically significant (Mann-Whitney U test; $U_s = 52.5$; $df = 20, 20$; $P < 0.002$).

The data in Table 3 are almost exclusively based on US sources (3 out of 20 references from Europe), whereas those in Table 2 are from both European and US sources (10 out of 20 references from Europe). The US studies are generally from more southern and warmer

Table 4.—Diet breadth (B) of five groups each of web-weaving spiders and hunting spiders; higher values indicate a less specialized feeding behavior (same data used as in Tables 2, 3).

Spider group	Diet breadth B	
	Mean \pm SE	Range
Web-weavers:		
Tetragnathidae	1.87 \pm 0.40	1.24–2.96
Araneidae	3.42 \pm 0.28	2.86–4.19
Theridiidae	3.20 \pm 0.60	1.70–4.52
Linyphiidae	2.55 \pm 0.30	1.85–3.20
Dictynidae	2.00 \pm 0.42	1.13–3.00
Overall mean	2.61 \pm 0.22	
Hunters:		
Oxyopidae (<i>Oxyopes</i>)	4.42 \pm 0.58	2.76–5.44
Oxyopidae (<i>Peucetia</i>)	4.34 \pm 0.34	3.42–4.86
Salticidae	4.38 \pm 0.33	3.45–4.89
Thomisidae	3.95 \pm 0.44	3.09–5.17
Lycosidae	3.90 \pm 0.69	2.65–5.58
Overall mean	4.20 \pm 0.20	

latitudes than the European ones (so far, most studies on the natural diets of hunters in crops available in the literature are from the southern US). Furthermore, the majority of US studies were conducted in structurally complex crops such as cotton and soybean fields, whereas most European studies were from cereal crops with a less complex (i.e., prevalently vertical) vegetation structure. Differences in geographic latitude as well as vegetation structure could influence the prey availabilities. Thus, the question arises whether the result of a greater diet breadth of the hunters observed in this study (Table 4) eventually is due to biases in the data set (the web and hunting spiders being studied in different crops and continents). To rule out this possibility, hunters and web-weavers should be analysed under comparable conditions (i.e., in the same field with identical prey availabilities).

Studies in which both hunters and web-weavers were evaluated in the same fields were published by Nyffeler (1982), Nyffeler & Sterling (1994), and Bardwell & Averill (1997). Based on these studies the diet breadth of web spiders and hunting spiders was assessed comparatively (Table 5). In Nyffeler's (1982) study in winter wheat fields near Zurich, Switzerland, hunters (represented by *Pardosa* spp. wolf spiders) had a greater diet

Table 5.—Diet breadth (*B*) of web-weaving spiders vs. hunting spiders in winter wheat, cotton, and cranberry, based on data from: ¹ Nyffeler (1982); ² Nyffeler & Sterling (1994); ³ Nyffeler et al. (1992a); ⁴ Bardwell & Averill (1997).

Crop	Foraging strategy	Spider species	Diet breadth <i>B</i>
WHEAT:	Web-weavers	<i>Tetragnatha extensa</i> ¹	1.24
		<i>Theridion impressum</i> ¹	2.90
		Erigoninae (pooled data) ¹	3.10
		<i>Achaearanea riparia</i> ¹	3.70
		Hunters	<i>Pardosa</i> spp. (pooled data) ¹
COTTON:	Web-weavers	<i>Tetragnatha laboriosa</i> ²	1.36
		<i>Latrodectus mactans</i> ²	1.70
		<i>Dictyna segregata</i> ²	2.37
		<i>Neoscona arabesca</i> ²	2.86
		<i>Acanthepeira stellata</i> ²	3.29
	Hunters	<i>Oxyopes salticus</i> ³	4.73
		<i>Oxyopes salticus</i> ²	4.76
CRANBERRY:	Web-weavers	(pooled data) ⁴	3.17
	Hunters	(pooled data) ⁴	4.69

breadth than the web-weavers (represented by orb weavers, sheet web-weavers, and tangle web-weavers) (Table 5). Likewise, in Texas cotton fields, the numerically dominant hunters (*Oxyopes salticus* and *Peucetia viridans*) exhibited greater diet breadth than several species of web-weavers (Table 5) (see Nyffeler et al. 1992a; Nyffeler & Sterling 1994). Furthermore, the data presented by Bardwell & Averill (1997) from cranberry bogs in Massachusetts suggest that the hunting spiders exhibited greater diet breadth than the web-weavers (pooled data for all hunters vs. web-weavers) (Table 5). Thus, in agroecosystems there seems to be a consistent trend of greater diet breadth of hunters compared to web-weavers regardless of crop type or geographic region investigated.

How do we explain this difference? Web spiders are stationary predators that wait for food to come to them (i.e., ‘sit-and-wait’ strategy). The prime requirement for the ‘sit-and-wait’ strategy is a food that moves (Turnbull 1973). A large proportion of web spiders spin aerial webs, with which they filter the aerial plankton (see Kajak 1965; Chacon & Eberhard 1980; Nentwig 1980). Others spin webs adapted to capture walking, crawling, or jumping prey (Turnbull 1973). Most web-weavers depend largely on relatively few prey groups available in high numbers in a particular environment (see Bristowe 1941; Turn-

bull 1960; Nyffeler & Benz 1979, Sunderland et al. 1986; Nentwig 1987; Alderweireldt 1994). In contrast, hunting spiders, by and large, seem to be less restricted in their diet (see Turnbull 1973). Representatives of various hunting spider families (e.g., Oxyopidae, Salticidae, Thomisidae, Lycosidae) have been reported to feed on both moving and motionless prey, which is indicative of a more mobile foraging strategy (see Nyffeler et al. 1990; Jackson & Tarsitano 1993). It is quite possible that the greater diet breadth of the hunting spiders (Table 4) simply reflects their greater opportunities to actively seek out suitable food due to their higher mobility (see Turnbull 1973).

There is observational evidence that hunting spiders can narrow their diet breadth significantly at times when a suitable prey type becomes locally superabundant relative to other prey (see Kiritani et al. 1972; Dean et al. 1987; Nyffeler et al. 1992b, 1994b). Thus, the greater diet breadth observed in the hunters (Table 4) does not necessarily imply that they require a more diverse diet than the web-weavers. It may instead show that they have a better chance of finding suitable food than web-weavers in agroecosystems (Young & Edwards 1990). However, there are exceptions to the rule (Turner & Polis 1979). Several members of the hunter families Thomisidae, Salticidae, Clubionidae, Gnaphosidae and Zo-

dariidae are known to specialize on ants (see Nentwig 1986, 1987).

Most spiders feed on prey that are small relative to their own size (prey length \leq spider length) (Wise 1993). Feeding experiments with a variety of spider species and a model prey (crickets) conducted in the laboratory revealed that the optimal prey length ranges from 50–80% of the spiders' own length (Nentwig 1987). Nentwig's laboratory data are fully supported by observations in the field (Hayes & Lockley 1990; Nyffeler et al. 1987b, c, 1992a). Overall, spider individuals of small size (including large percentages of immatures) numerically dominate the faunas of field crops, and these feed primarily on tiny prey organisms (< 4 mm in length) (LeSar & Unzicker 1978; Young & Edwards 1990; Nyffeler et al. 1994a).

ACKNOWLEDGMENTS

I extend my gratitude to Matt Greenstone, Wolfgang Nentwig, Keith Sunderland, Soeren Toft, and an anonymous reviewer for their comments on earlier drafts of this paper.

LITERATURE CITED

- Agnew, C.W. & J.W. Smith, Jr. 1989. Ecology of spiders (Araneae) in a peanut agroecosystem. *Environ. Entomol.*, 18:30–42.
- Alderweireldt, M. 1994. Prey selection and prey capture strategies of linyphiid spiders in high-input agricultural fields. *Bull. British Arachnol. Soc.*, 9:300–308.
- Bardwell, C.J. & A.L. Averill. 1997. Spiders and their prey in Massachusetts cranberry bogs. *J. Arachnol.*, 25:31–41.
- Bristowe, W.S. 1941. *The Comity of Spiders*. Ray Society, London.
- Chacon, P. & W.G. Eberhard. 1980. Factors affecting numbers and kinds of prey caught in artificial spider webs, with considerations of how orb webs trap prey. *Bull. British Arachnol. Soc.*, 5:29–38.
- Colwell, R.K. & D.J. Futuyma. 1971. On the measurement of niche breadth and overlap. *Ecology*, 52:567–576.
- Culin, J.D. & K.V. Yeorgan. 1982. Feeding behavior and prey of *Neoscona arabesca* (Araneae: Araneidae) and *Tetragnatha laboriosa* (Araneae: Tetragnathidae) in soybean fields. *Entomophaga*, 27:417–424.
- Dean, D.A., W.L. Sterling, M. Nyffeler & R.G. Breene. 1987. Foraging by selected spider predators on the cotton fleahopper and other prey. *Southwest. Entomol.*, 12:263–270.
- Greenstone, M.H. 1979. Spider feeding behaviour optimises dietary essential amino acid composition. *Nature*, 282:501–503.
- Hallander, H. 1970. Prey, cannibalism and microhabitat selection in the wolf spiders *Pardosa chelata* (O.E. Müller) and *P. pullata* (Clerck). *Oikos*, 21:337–340.
- Hassell, M.P. 1978. *The Dynamics of Arthropod Predator–prey Systems*. Monographs in Population Biology 13. Princeton Univ. Press, New Jersey.
- Hayes, J.L. & T.C. Lockley. 1990. Prey and nocturnal activity of wolf spiders (Araneae: Lycosidae) in cotton fields in the Delta region of Mississippi. *Environ. Entomol.*, 19:1512–1518.
- Heidger, C. & W. Nentwig. 1986. The prey of *Dictyna arundinacea* (Araneae: Dictynidae). *Zool. Beitr. (NF)*, 29:185–192.
- Heidger, C. & W. Nentwig. 1989. Augmentation of beneficial arthropods by strip-management. 3. Artificial introduction of a spider species which preys on wheat pest insects. *Entomophaga*, 34: 511–522.
- Jackson, R.R. 1977. Prey of the jumping spider *Phidippus johnsoni* (Araneae: Salticidae). *J. Arachnol.*, 5:145–149.
- Jackson, R.R. 1992. Eight-legged tricksters (Spiders that specialize in catching other spiders). *BioScience*, 42:590–598.
- Jackson, R.R. & M.S. Tarsitano. 1993. Responses of jumping spiders to motionless prey. *Bull. British Arachnol. Soc.*, 9:105–109.
- Kajak, A. 1965. An analysis of food relations between the spiders —*Araneus cornutus* Clerck and *Araneus quadratus* Clerck — and their prey in meadows. *Ekol. Polska (A)*, 13:717–764.
- Kiritani, K., S. Kawahara, T. Sasaba & F. Nakasuji. 1972. Quantitative evaluation of predation by spiders on the green rice leafhopper, *Nephotettix cincticeps* Uhler, by a sight-count method. *Res. Popul. Ecol.*, 13:187–200.
- LeSar, C.D. & J.D. Unzicker. 1978. Life history, habits, and prey preferences of *Tetragnatha laboriosa* (Araneae: Tetragnathidae). *Environ. Entomol.*, 7:879–884.
- Levins, R. 1968. Evolution in changing environments: Some theoretical explorations. Monographs in Population Biology. 2. Princeton Univ. Press, New Jersey.
- Lockley, T.C. & O.P. Young. 1987. Prey of the striped lynx spider, *Oxyopes salticus* (Araneae, Oxyopidae), on cotton in the Delta area of Mississippi. *J. Arachnol.*, 14:395–397.
- Morse, D.H. 1983. Foraging patterns and time budgets of the crab spiders *Xysticus emertoni* Keyserling and *Misumena vatia* (Clerck) (Araneae, Thomisidae) on flowers. *J. Arachnol.*, 11:87–94.
- Nentwig, W. 1980. The selective prey of linyphiid-like spiders and of their space webs. *Oecologia*, 45:236–243.

- Nentwig, W. 1986. Non-webbuilding spiders: prey specialists or generalists? *Oecologia*, 69:571–576.
- Nentwig, W. 1987. The prey of spiders. Pp. 249–263. *In* *Ecophysiology of Spiders*. (W. Nentwig, ed.). Springer-Verlag, Berlin, New York.
- Nyffeler, M. 1982. Field studies on the ecological role of the spiders as insect predators in agroecosystems. Ph.D. Dissertation, Swiss Fed. Inst. Technol., Zurich, 174 pp.
- Nyffeler, M. & G. Benz. 1979. Studies on the ecological importance of spider populations for the vegetation of cereal and rape fields. *J. Appl. Entomol.*, 87:348–376 (in German).
- Nyffeler, M. & G. Benz. 1988a. Prey analysis of the spider *Achaearanea riparia* (Blackw.) (Araneae, Theridiidae), a generalist predator in winter wheat fields. *J. Appl. Entomol.*, 106:425–431.
- Nyffeler, M. & G. Benz. 1988b. Prey and predatory importance of micryphantid spiders in winter wheat fields and hay meadows. *J. Appl. Entomol.*, 105:190–197.
- Nyffeler, M. & G. Benz. 1988c. Feeding ecology and predatory importance of wolf spiders (*Paradosa* spp.) (Araneae, Lycosidae) in winter wheat fields. *J. Appl. Entomol.*, 106:123–134.
- Nyffeler, M. & R.G. Breene. 1990a. Spiders associated with selected European hay meadows and the effects of habitat disturbance, with the predation ecology of the crab spiders, *Xysticus* spp. (Araneae: Thomisidae). *J. Appl. Entomol.*, 110:149–159.
- Nyffeler, M. & R.G. Breene. 1990b. Evidence of low daily food consumption by wolf spiders in meadowland and comparison with other cursorial hunters. *J. Appl. Entomol.*, 110:73–81.
- Nyffeler, M. & W.L. Sterling. 1994. Comparison of the feeding niche of polyphagous insectivores (Araneae) in a Texas cotton plantation: Estimates of niche breadth and overlap. *Environ. Entomol.*, 23:1294–1303.
- Nyffeler, M., R.G. Breene, D.A. Dean & W.L. Sterling. 1990. Spiders as predators of arthropod eggs. *J. Appl. Entomol.*, 109:490–501.
- Nyffeler, M., D.A. Dean & W.L. Sterling. 1987a. Feeding ecology of the orb-weaving spider *Argiope aurantia* (Araneae: Araneidae), in a cotton agroecosystem. *Entomophaga*, 32:367–375.
- Nyffeler, M., D.A. Dean & W.L. Sterling. 1987b. Evaluation of the importance of the striped lynx spider, *Oxyopes salticus* (Araneae: Oxyopidae), as a predator in Texas cotton. *Environ. Entomol.*, 16:1114–1123.
- Nyffeler, M., D.A. Dean & W.L. Sterling. 1987c. Predation by green lynx spider, *Peucetia viridans* (Araneae: Oxyopidae), inhabiting cotton and woolly croton plants in East Texas. *Environ. Entomol.*, 16:355–359.
- Nyffeler, M., D.A. Dean & W.L. Sterling. 1988a. The southern black widow spider, *Latrodectus mactans* (Araneae, Theridiidae), as a predator of the red imported fire ant, *Solenopsis invicta* (Hymenoptera, Formicidae), in Texas cotton fields. *J. Appl. Entomol.*, 106:52–57.
- Nyffeler, M., D.A. Dean & W.L. Sterling. 1988b. Prey records of the web-building spiders *Dictyna segregata* (Dictynidae), *Theridion australe* (Theridiidae), *Tidarren haemorrhoidale* (Theridiidae), and *Frontinella pyramitela* (Linyphiidae) in a cotton agroecosystem. *Southwest. Nat.*, 33:215–218.
- Nyffeler, M., D.A. Dean & W.L. Sterling. 1989. Prey selection and predatory importance of orb-weaving spiders (Araneae: Araneidae, Uloboridae) in Texas cotton. *Environ. Entomol.*, 18:373–380.
- Nyffeler, M., D.A. Dean & W.L. Sterling. 1992a. Diets, feeding specialization, and predatory role of two lynx spiders, *Oxyopes salticus* and *Peucetia viridans* (Araneae: Oxyopidae), in a Texas cotton agroecosystem. *Environ. Entomol.*, 21:1457–1465.
- Nyffeler, M., W.L. Sterling & D.A. Dean. 1992b. Impact of the striped lynx spider (Araneae: Oxyopidae) and other natural enemies on the cotton fleahopper *Pseudatomoscelis seriatus* (Hemiptera: Miridae) in Texas cotton. *Environ. Entomol.*, 21:1178–1188.
- Nyffeler, M., W.L. Sterling & D.A. Dean. 1994a. Insectivorous activities of spiders in United States field crops. *J. Appl. Entomol.*, 118:113–128.
- Nyffeler, M., W.L. Sterling & D.A. Dean. 1994b. How spiders make a living. *Environ. Entomol.*, 23:1357–1367.
- Randall, J.B. 1982. Prey records of the green lynx spider, *Peucetia viridans* (Hentz) (Araneae, Oxyopidae). *J. Arachnol.*, 10:19–22.
- Riechert, S.E. & L. Bishop. 1990. Prey control by an assemblage of generalist predators: spiders in garden test systems. *Ecology*, 71:1441–1450.
- Riechert, S.E. & J.M. Harp. 1987. Nutritional ecology of spiders. Pp. 645–672. *In* *Nutritional Ecology of Insects, Mites, and Spiders*. (F. Slansky & J.G. Rodriguez, eds.). John Wiley, New York.
- Riechert, S.E. & J. Luczak. 1982. Spider foraging: behavioral responses to prey. Pp. 353–385. *In* *Spider Communication: Mechanisms and Ecological Significance*. (P.N. Witt & J.S. Rovner, eds.). Princeton Univ. Press, New Jersey.
- Rypstra, A.L. 1982. Building a better insect trap; an experimental investigation of prey capture in a variety of spider webs. *Oecologia*, 52:31–36.
- Sunderland, K.D., A.M. Fraser & A.F.G. Dixon. 1986. Distribution of linyphiid spiders in relation to capture of prey in cereal fields. *Pedobiologia*, 29:367–375.
- Toft, S. 1995. Value of the aphid *Rhopalosiphum*

- padi* as food for cereal spiders. J. Appl. Ecol., 32:552–560.
- Turnbull, A.L. 1960. The prey of the spider *Linyphia triangularis* (Clerck) (Araneae, Linyphiidae). Canadian J. Zool., 38:859–873.
- Turnbull, A.L. 1973. Ecology of the true spiders (Araneomorphae). Annu. Rev. Entomol., 18:305–348.
- Turner, M. 1979. Diet and feeding phenology of the green lynx spider, *Peucetia viridans* (Araneae: Oxyopidae). J. Arachnol., 7:149–154.
- Turner, M. & G.A. Polis. 1979. Patterns of co-existence in a guild of raptorial spiders. J. Anim. Ecol., 48:509–520.
- Uetz, G.W., J. Bischoff & J. Raver. 1992. Survivorship of wolf spiders (Lycosidae) reared on different diets. J. Arachnol., 20:207–211.
- Uetz, G.W., A.D. Johnson & D.W. Schemske. 1978. Web placement, web structure, and prey capture in orb-weaving spiders. Bull. British Arachnol. Soc., 4:141–148.
- Wise, D.H. 1993. Spiders in Ecological Webs. Cambridge Univ. Press, Cambridge, U.K.
- Yeargan, K.V. 1975. Prey and periodicity of *Paradosa ramulosa* (McCook) in alfalfa. Environ. Entomol., 4:137–141.
- Young, O.P. 1989. Field observations of predation by *Phidippus audax* (Araneae: Salticidae) on arthropods associated with cotton. J. Entomol. Sci., 24:266–273.
- Young, O.P. & G.B. Edwards. 1990. Spiders in United States field crops and their potential effect on crop pests. J. Arachnol., 18:1–27.

Manuscript received 1 May 1998, revised 1 October 1998.